

Mauro H. Chagas/Dietmar Schmidtbleicher

Auswirkungen von Beweglichkeitstraining auf die Bewegungsamplitude, Dehnungsspannung und Dehngrenze nach einer Trainings- und Detrainingsperiode*

Literatur

- Akeson, W.H.; Amiel, D. & Woo, S.L-Y. (1980). Immobility effects on synovial joints the pathomechanics of joint contraction. *Biorheology*, 17, 95-110.
- Alter, M. J. (1996). *Science of flexibility*. Champaign (2nd edition): Human Kinetics.
- Bandy, W. & Irion, M. (1994). The effect of time on static stretch on the flexibility of the hamstring muscles. *Phys. Ther.*, 74, 485-850.
- Bandy, W. & Irion, M. (1998). The effect of static stretch and dynamic range of motion training on the flexibility of the hamstring muscles. *J. Orthop. Sports Phys. Ther.*, 27, 295-300.
- Bavel, H.V., Drost, M.R., Wielders, J.D.L., Huyghe, J.M., Huisson, A. & Janssen, J.D. (1996). Strain distribution on rat medial gastrocnemius (MG) during passive stretch. *Journal of Biomechanics*, 29, 1069-1074.
- Billeter, R. & Hoppeler, H. (1992): Muscular basis of strength. In P.V. Komi (Ed.), *Strength and Power in Sport* (pp. 39-63). London: Blackwell Scientific Publications.
- Borg, T.K. & Caulfield, J.B. (1980). Morphology of connective tissue in skeletal muscle. *Tissue & Cell*, 12, 197-207.
- Chagas, M.H. & Schmidtbleicher, D. (2000). *Effects of two stretching methods on the range of motion, passive muscle tension and retention of flexibility*. Proceedings of the 5th Annual Congress of the European College of Sport Science. Jyväskylä, 201.
- Chagas, M.H. & Schmidtbleicher, D. (2001a). *Improvement of the range of motion and muscle passive tension during and after long-term stretching training*. Proceedings of the 6th Annual Congress of the European College of Sport Science. Cologne, 788.
- Chagas, M.H. & Schmidtbleicher, D. (2001b). *Effect of increase of the range of motion on muscle strength performance*. Proceedings of the 6th Annual Congress of the European College of Sport Science. Cologne, 1040.
- Crawford, G.N.C. (1973). The growth of striated muscle immobilized in extension. *J. Anatomy*, 114, 165-183.
- Culav, E.M., Clark, C.H. & Merrilees, M.J. (1999). Connective tissues: matrix composition and its relevance to physical therapy. *Phys. Ther.*, 79, 308-319.
- Curwin, S.L., Vailas, A.C. & Wood, J. (1988). Immature tendon adaptation to strenuous exercise. *J. Appl. Physiol.*, 65, 2297-2301.
- De Deyne, P.G. (2001). Application of passive stretch and its implications for muscle fibers. *Phys. Ther.*, 81, 819-827.
- Erickson, H.P. (1997). Stretching single protein molecules: Titin is a weird spring. *Science*, 276, 1090-1092.
- Etnyre, B.R. & Abraham, L.D. (1986). Gains in range of ankle dorsiflexion using three popular stretching techniques. *Am. J. Phys. Med.*, 65, 189-196.
- Etnyre, B.R. & Lee, E.J. (1988). Chronic and acute flexibility of men and women using three different stretching techniques. *Res. Quart.*, 59, 222-228.
- Eyre, D.R. & Oguchi, H. (1980). The hydroxyproline cross-links of skeletal collagens: their measurement, properties and a proposed pathway of formation. *Biochem. Biophys. Res. Commun.*, 92, 403-410.
- Freiwald, J. & Engelhardt, M. (1994). Beweglichkeit und ihre Einschränkungen: vor Training und Therapie Faktoren genau analysieren. *TW Sport + Medizin*, 6, 327-336.
- Fürst, D.O. (1999). Titin, ein molekularer Gigant regiert im quergestreiften Muskel. *Deutsche Zeitschrift für Sportmedizin*, 50, 218-222.
- Gajdosik, R.L. (1991). Effects of static stretching on the maximal length and resistance to passive stretch of short hamstring muscles. *J. Orthop. Sports Phys. Ther.*, 14, 250-255.
- Gajdosik, R.L. (1995). Flexibility or muscle length? [letter; comment]. *Phys. Ther.*, 75, 238-239.
- Gajdosik, R.L. (1997). Influence of age on calf muscle length and passive stiffness variables at different stretch velocities. *Isokinetics and Exercise Science*, 6, 163-174.
- Gajdosik, R.L. (2001). Passive extensibility of skeletal muscle: review of the literature with clinical implications. *Clinical Biomechanics*, 16, 87-101.
- Göeken, L.N. & Hof, L. (1994). Instrumental straight-leg raising: results in patients. *Arch. Phys. Med. Rehabil.*, 75, 470-477.
- Goldspink, G. (1992). Cellular and molecular aspects of adaptation in skeletal muscle. In P.V. Komi (Ed.), *Strength and power in sport* (pp. 211-229). London: Blackwell Scientific Publications.
- Halbertsma, J.P.K. & Göeken, L.N.H. (1994). Stretching exercises: effect on passive extensibility and stiffness in short hamstrings of healthy subjects. *Arch. Phys. Med. Rehabil.*, 75, 976-981.
- Halbertsma, J.P.K., Mulder, I., Göeken, L.N.H. & Eisma, W.H. (1999). Repeated passive stretching: acute effect on the passive muscle moment and extensibility of short hamstrings. *Archives of physical medicine and rehabilitation*, 80, 407-414.
- Handel, M., Horstmann, T. & Dickhuth, H.-H. (1997). Effects of contract-relax stretching training on muscle performance in athletes. *Eur. J. Appl. Physiol.*, 76, 400-408.
- Horowitz, R. (1992). Passive force generation and titin isoforms in mammalian skeletal muscle. *Biophys. J.*, 61, 392-398.
- Hortobágyi, J., Faludi, J., Tihanyi, J. & Merkely, B. (1985). Effects of intense „stretching“ flexibility training on the mechanical profile of the knee extensors and on the range of motion of the hip joint. *Int. J. Sports Med.*, 6, 317-321.
- Huijing, P.A. (1992). Mechanical muscle models. In P.V. Komi (Ed.), *Strength and Power in Sport* (pp. 130-150). London: Blackwell Scientific Publications.
- Huijing, P.A. (1997). *Muscle as collagen fiber reinforced composite material: force transmission in muscle and whole limbs*. Proceedings of the XVIth ISB Congress. Tokio, 7.
- Hutton, R.S. (1992). Neuromuscular basis of stretching exercises. In P.V. Komi (Ed.), *Strength and power in sport* (pp. 29-38). London: Blackwell Scientific Publications.
- Kibele, A. (1998). Maximalkraftzuwachs ohne Leistungsgewinn. *Leistungssport*, 28 (3), 45-49.
- Kovanen, V.H., Suominen, H. (1989). Age- and training-related changes in the collagen metabolism of rat skeletal muscle. *Eur. J. Appl. Physiol.*, 58, 765-771.
- Kubo, K., Kanehisa, H., Kawakami, Y. & Fukunaga, T. (2002a). Effect of stretching training on the viscoelastic properties of human tendon structures in vivo. *J. Appl. Physiol.*, 92, 595-601.
- Kubo, K., Kanehisa, H. & Fukunaga, T. (2002b). Effects of resistance and stretching training programmes on the viscoelastic properties of human tendon structures in vivo. *J. Physiol.*, 538, 219-226.
- Latash, M. L. & Zatsiorsky, V. M. (1993). Joint stiffness: myth or reality? *Hum. Mov. Sci.*, 12, 653-692.
- Magnusson, S.P., Simonsen, E.B., Aagaard, P., Gleim, G.W., McHugh, M.P. & Kjaer, M. (1995). Viscoelastic response to repeated static stretching in human skeletal muscle. *Scand. J. Med. Sci. Sports*, 5, 342-347.
- Magnusson, S.P., Simonsen, E.B., Aagaard, P., Sørensen, H. & Kjaer, M. (1996a). A mechanism for altered flexibility in human skeletal muscle. *J. Physiol.*, 497, 291-298.
- Magnusson, S.P., Simonsen, E.B., Aagaard, P. & Kjaer, M. (1996b). Biomechanical responses to repeated stretches in human hamstring muscle in vivo. *Am. J. Sports Med.*, 24, 622-628.
- Magnusson, S.P., Aagaard, P., Larsson, B. & Kjaer, M. (2000). Passive energy absorption by human muscle-tendon unit is unaffected by increase in intramuscular temperature. *J. Appl. Physiol.*, 88, 1215-1220.
- Marchettini, P. (1993). Muscle pain: animal and human experimental and clinical studies. *Muscle & Nerve*, 16, 1033-1039.
- Marschall, F. (1999). Wie beeinflussen unterschiedliche Dehnintensitäten kurzfristig die Veränderung der Bewegungsreichweite. *Deutsche Zeitschrift für Sportmedizin*, 50, 5-9.
- McComas, A. J. (1996). *Skeletal muscle: form and function*. Champaign: Human Kinetics.
- McHugh, M.P., Kremenik, I.J., Box, M.B. & Gleim, G. W. (1998). The role of mechanical and neural restraints to joint range of motion during passive stretch. *Med. Sci. Sports Exer.*, 30, 928-932.
- Mense, S. (1984). Muskelrezeptoren mit nociception Eigenschaften. In M. Berger, F. Gerstenbrand & K. Lewit (Hrsg.), *Schmerz und Bewegungssystem* (S. 17-26). Stuttgart: Gustav Fischer.
- Mense, S. (1988). Verhalten von Nozizeptoren im normalen und entzündeten Muskel. In R. Spintge & R. Droh (Hrsg.), *Schmerz und Sport* (S. 199-206). Berlin: Springer Verlag
- Mense, S. (1993): Nociception from muscle in relation to clinical muscle pain. *Pain*, 54, 241-289.
- Meßlinger, K. (1997). Was ist ein Nozizeptor? *Der Anaesthetist*, 46, 142-153.
- Mohr, K.J., Pink, M.M., Elsner, C. & Kvitne, R.S. (1998) Electromyographic investigation of stretching: the effect of warm-up. *Clinical J. of Sports Med.*, 8, 215-220.
- Moore, M.A. & Hutton, R.S. (1980). Electromyographic investigation of muscle stretching techniques. *Med. Sci. Sports Exer.*, 12, 322-329.
- Ott, H. (2000). *Entwicklung und Erprobung eines diagnostischen Untersuchungsverfahrens für die Beweglichkeit der ischiocruralen Muskulatur*. Dissertation, Frankfurt.
- Purslow, R.P. (1989). Strain-induced reorientation of an intramuscular connective tissue network: implications for passive muscle elasticity. *J. Biomechanics*, 22, 21-31.
- Rauber, A. & Kopsch, F. (1987). *Anatomie des Menschen*. Band I. Stuttgart: Thieme.

- Rowe, R. W. (1981). Morphology of perimysial and endomy-sial connective tissue in skeletal muscle. *Tissue & Cell*, 13, 681-690.
- Roy, R.R. & Edgerton, V.R. (1992). Skeletal architecture and performance. In P.V. Komi (Ed.), *Strength and Power in Sport* (pp. 115-129). London: Blackwell Scientific Publications.
- Sady, S.P., Wartman, M. & Blanke, D. (1982). Flexibility training: ballistic, static or proprioceptive neuromuscular facilitation? *Arch. Phys. Med. Rehabil.*, 63, 261-263.
- Sale, D.G. (1991). Testing strength and power. In J.D. Mac-Dougall, H.A. Wenger & H.J. Green (Eds.), *Physiological testing of the high-performance athlete* (pp. 21-106). Cham-paign: Human Kinetics.
- Schönthaler, S.R., Ohlendorf, K., Ott, H., Meyer, T., Kinder-mann, W. & Schmidbleicher, D. (1998). Biomechanische und neurophysiologische Parameter zur Erfassung der Dehnbar-keit von Muskel-Sehnen-Einheiten. *Deutsche Zeitschrift für Sportmedizin*, 49, 223-230.
- Schönthaler, S.R. & Ohlendorf, K. (2000). *Biomechanische und neurophysiologische Veränderungen nach ein- und mehrfachseriellem passiv-statischem Beweglichkeitstrai-ning*. Dissertation, Frankfurt/Main.
- Shadwick, R.E. (1990). Elastic energy storage in tendons: mechanical differences related to function and age. *J. Appl. Physiol.*, 68, 1033-1040.
- Simonsen, E.B., Klitgaard, H. & Bojsen-Moller, F. (1995). The influence of strength training, swim training and ageing on the achilles tendon and m. soleus of the rat. *J. Sports Sci.*, 13, 291-295.
- Smith L.L., Brunetz, M.H., Chenier, T.C., McCammon, M.R., Houmard, J.A., Franklin, M.E. & Israel R.G. (1993). The ef-fects of static and ballistic stretching on delayed onset mus-cle soreness and creatine kinase. *Res. Quart.*, 64, 103-107.
- Stacey, M.J. (1969). Free nerve endings in skeletal muscle of the cat. *J. Anatomy*, 105, 231-254.
- Stone, M.H. (1992). Connective tissue and bone response to strength training. In P.V. Komi (Ed.), *Strength and Power in Sport* (pp. 279-290). London: Blackwell Scientific Publicati-ons.
- Sullivan, M.K., DeJulia, J.J. & Worrel, T.W. (1992). Effect of pelvic position and stretching method on hamstring muscle flexibility. *Med. Sci. Sports Exerc.*, 24, 1383-1389.
- Suominen, H., Heikkinen, E. & Parkatti, T. (1977). Effect of eight weeks' physical training on muscle and connective tis-sue of the m. vastus lateralis in 69-year-old men and women. *J. Gerontology*, 32, 33-37.
- Tabary, J.C., Tabary, C., Tardieu, G. & Goldspink, G. (1972). Physiology and structural changes in the cat's soleus muscle due to immobilization at different lengths by plaster casts. *J. Physiol.*, 224, 231-244.
- Tittel, K. (1990). Beschreibende und funktionelle Anatomie des Menschen (11. Auflage). Stuttgart: Gustav Fischer.
- Toft, E., Espersen, G.T., Kalund, S., Sinkjaer, T. & Hornemann, B.C. (1989). Passive tension of the ankle before and after stretching. *Am. J. Sports Med.*, 17, 489-494.
- Trinick, J. (1991). Elastic filaments and giant proteins in mus-cle. *Curr. Opin. Cell Biol.*, 3, 112-118.
- Trotter, J.A. & Purslow, P.P. (1992). Functional morphology of the endomysium in series fibered muscles. *J. Morphology*, 212, 109-122.
- Ullrich, K. & Gollhofer, A. (1994). Physiologische Aspekte und Effektivität unterschiedlicher Dehnmethoden. *Deutsche Zeit-schrift für Sportmedizin*, 45, 336-345.
- Viidik, A. (1973). Functional properties of collagenous tis-sues. In D.A. Hall & D.S. Jackson (Eds.), *International review of connective tissue research* (pp. 127-209). London: Acade-mic Press.
- Viidik, A. (1980). Elastomechanik biologischer Gewebe. In H. Cotta. (Hrsg.): *Die Belastungstoleranz des Bewegungsappa-rates* (S. 124-136). Stuttgart: Georg Thieme Verlag.
- Viidik, A. (1986). Adaptability of connective tissue. In: B. Sal-tin (Ed.), *Biochemistry of exercise* (pp. 545-562). Cham-paign: Human Kinetics.
- Wallin, D., Ekblom, B., Grahn, R. & Nordenborg, T. (1985): Improvement of muscle flexibility: a comparison between two techniques. *Am. J. Sports Med.*, 13, 263-268.
- Wang, K.J., McCarter, R., Wright, J., Beverly, J. & Ramirez-Mitchell, R. (1991): Regulation of skeletal muscle stiffness and elasticity by titin isoforms: a test of the segmental ex-tension model of resting tension. *Proc. Natl. Acad. Sci. USA*, 88, 7101-7105.
- Wang, K.J., McCarter, R., Wright, J., Beverly, J. & Ramirez-Mitchell, R. (1993). Viscoelasticity of the sarcomere matrix of skeletal muscles. *Biophys. J.*, 64, 1161-1177.
- Wiemann, K. (1991). Beeinflussung muskulärer Parameter durch ein zehnwöchiges Dehnungstraining. *Sportwissen-schaft*, 21, 295-305.
- Wiemann, K. & Leisner, S. (1996). Extreme Hüftbeugefähig-keit von Turnern: Sind längere ischiocrurale Muskeln die Ur-sache? *TW Sport + Medizin*, 8, 103-108.
- Wiemann, K., Klee, A. & Stratmann, M. (1998). Filamentäre Quellen der Muskel-Ruhespannung und die Behandlung mus-kulärer Dysbalancen. *Deutsche Zeitschrift für Sportmedizin*, 49, 111-118.
- Wiemann, K. & Klee, A. (2000). Die Bedeutung von Dehnen und Stretching in der Aufwärmphase vor Höchstleistungen. *Leistungssport*, 30 (4), 5-9.
- Williams, P.E. & Goldspink, G. (1978). Changes in sarcomere length and physiological properties in immobilized muscle. *J. Anatomy*, 127, 459 - 468.
- Woo, S. L-Y., Matthews, J.V., Akeson, W.H., Amiel, D. & Con-very, F.R. (1975). Connective tissue response to immobility. *Arthritis and Rheumatism*, 18, 257-264.
- Woo, S. L-Y., Mark, A., Gomes, B.A., Woo, Y-K. & Akeson, W.H. (1982). Mechanical properties of tendons and ligaments II: the relationships of immobilization and exercise on tissue re-modeling. *Biorheology*, 19, 397-408.
- Wydra, G., Bös, K. & Karisch, G. (1991). Zur Effektivität ver-schiedener Dehntechniken. *Deutsche Zeitschrift für Sport-medizin*, 42, 386-534.
- Youdas, J.W., Krause, D.A., Egan, K.S., Therneau, T.M. & Las-kowski, E.R. (2003). The effect of static stretching of the calf muscle-tendon unit on active ankle dorsiflexion range of mo-tion. *J. Orthop. Sports Phys. Ther.*, 33, 295-300.
- Zachazewski, J.E. (1990). Flexibility for sports. In B. Sanders (Ed.), *Sport physical therapy* (pp. 201-238). Norwalk: Applet-on & Lange.
- Zernicke, R.F. & Loitz, B.J. (1992). Exercise-related adapta-tions in connective tissue. In P.V. Komi (Hrsg.), *Strength and power in sport* (pp. 77-95). London: Blackwell Scientific Pu-blications.

*