

Salah A. Gahlul/Peter Hofmann

# SPRINT-INTERVALLTRAINING BEI JUGENDLICHEN FUSSBALLERN

## Auswirkungen auf die Verbesserung der aeroben und anaeroben Ausdauer-Leistungsfähigkeit

### Literatur

- Bailey, R. C., Olson, J., Pepper, S. L., Porszasz, J., Barstow, T. J. et al. (1995). The level and tempo of children's physical activities: an observational study. *Med. Sci. Sports Exerc.*, 27 (7), 1033-1041.
- Bangsbo, J., Nørregaard, L. & Thorsøe, F. (1991). Activity profile of competition soccer. *Can. J. Sports Sci.*, 16 (2), 110-116.
- Bangsbo, J. (1993). *The Physiology of Soccer – with Special Reference to Intense Intermittent Exercise*. August Krogh Institute, University of Copenhagen.
- Bangsbo, J. (1994). Physiological demands. In B. Ekblom (ed.), *Football (soccer)* (pp. 43-59). London: Blackwell.
- Bangsbo, J., Iaia, F. M. & Krstrup, P. (2008). The Yo-Yo Intermittent Recovery Test: A useful tool for evaluation of physical performance in intermittent sports. *Sports Medicine*, 38 (1), 37-51.
- Baquet, G., Berthoin, S., Dupont, G., Blondel, N., Fabre, C. et al. (2002). Effects of high intensity intermittent training on peak  $\text{VO}_2$  in prepubertal children. *Int. J. Sports Med.*, 23 (6), 439-444.
- Barkley, J. E., Epstein, L. H. & Roemmich, J. N. (2009). Reinforcing value of interval and continuous physical activity in children. *Physiology and Behavior*, 98, 31-36.
- Bickham, D. C., Bentley, D. J., Le Rossignol, P. F. & Cameron-Smith, D. (2006). The effects of short-term sprint training on MCT expression in moderately endurance trained runners. *Eur. J. Appl. Physiol.*, 96, 636-643.
- Conconi, F., Ferrari, M., Ziglio, P. G., Droghetti, P. & Codeca, L. (1982). Determination of the anaerobic threshold by a noninvasive field test in runners. *Journal of Applied Physiology*, 52, 869-873.
- Cureton, T. K. (1970). *Physical Workbook for Adult (Band I)*. Champaign (IL): Stipes Publishing.
- Dudley, G. A. & Djamil, R. (1985). Incompatibility of endurance- and strength-training modes of exercise. *J. Appl. Physiol.*, 59, 1446-1451.
- Dupont, G., Mathieu, D., Laurent, B., Nicolas, B., Wassim, M. et al. (2008). Yo-Yo intermittent recovery test versus the Université de Montréal Track Test: Relation with a high-intensity intermittent exercise. *J. Sci. Med. Sport*. doi:10.1016/j.jksam.2008/10.007.
- Dupont, G. & Berthoin, S. (2004). Time spent at high percentage of  $\text{VO}_{2\text{max}}$  for short intermittent runs: active versus passive recovery. *Can. J. Appl. Physiol.*, 29 (Suppl.), S2-S16.
- Epstein, L. H. K. C., Consalvi, A. R. & Paluch, R. A. (1999). Reinforcing value of physical activity as a determinant of child activity level. *Health Psychol.*, 18 (6), 599-603.
- Esfarjani, F. & Laursen, P. B. (2007). Manipulating high-intensity interval training: effects on  $\text{VO}_{2\text{max}}$ , the lactate threshold and 3000 m running performance in moderately trained males. *J. Sci. Med. Sport*, 10, 27-35.
- Ferrari, B. D., Impellizzeri, F. M., Rampinini, E., Castagna, C., Bishop, D. et al. (2008). Sprint vs. interval training in football. *Int. J. Sports Med.*, 29 (8), 668-674.
- Franke, J. (2011). *Ausdauertraining in ausgewählten Sportarten – extensives Dauerlauftraining versus intensives Intervaltraining in der Saisonvorbereitung*. Fakultät für Sportwissenschaft der Ruhr-Universität Bochum.
- Gamelin, F.-X., Baquet, G., Berthoin, S., Thevenet, D. et al. (2009). Effect of high intensity intermittent training on heart rate variability in prepubescent children. *Eur. J. Appl. Physiol.*, 105 (5), 731-738.
- Gibala, M. J., Little, J. P., van Essen, M., Wilkin, G. P., Burgomaster, K. A. et al. (2006). Short-term sprint interval versus traditional endurance training: similar initial adaptations in human skeletal muscle and exercise performance. *J. Physiol.*, 575, 901-911.
- Gunnarsson, T. P., Christensen, P. M., Holse, K., Christiansen, D. & Bangsbo, J. (2012). Effect of additional speed endurance training on performance and muscle adaptations. *Med. Sci. Sports Exerc.*, 44 (10), 1942-1948.
- Hazell, T. J., MacPherson, R. E. K., Gravelle, B. M. R. & Lemon, P. W. R. (2010). 10- or 30-s sprint interval training bouts enhance both aerobic and anaerobic performance. *Eur. J. Appl. Physiol.* doi: 10.1007/s00421-010-1474-y.
- Helgerud, J., Engen, L. C., Wisloff, U. & Hoff, J. (2001). Aerobic endurance training improves soccer performance. *Med. Sci. Sports Exerc.*, 33, 1925-1931.
- Helgerud, J., Hoydal, K., Wang, E., Karlsen, T., Berg, P. et al. (2007). Aerobic high-intensity intervals improve  $\text{VO}_{2\text{max}}$  more than moderate training. *Med. Sci. Sports Exerc.*, 39 (4), 665-671.
- Hill-Haas, S. V., Coutts, A. J., Rowsell, G. J. & Dawson, B. T. (2009). Generic versus small-sided game training in soccer. *Int. J. Sports Med.* doi:10.1055/s-0029-1220730.
- Hoff, J. & Helgerud, J. (2005). Endurance and strength training for soccer players – physiological considerations. *Sports Med.*, 34 (3), 165-180.
- Hofmann, P. & Gaisl, G. (1988). Der Conconi-Test und seine praktischen Anwendungen. *Condition*, 19 (7), 18-19.
- Hofmann, P., Gaisl, B., Stockinger, R. & Leitner, H. (1989). Modifikationen des Conconi-Tests für die Anwendung in Hallenspielsportarten. *Leistungssport*, 19 (3), 27 f.
- Hofmann, P. & Pokan, R. (2010). Value of the application of heart rate performance curve in sport. *Int. J. Sport Physiol. Perform.*, 5 (4), 437-447.
- Iaia, F. M., Hellsten, Y., Nielsen, J. J., Fernstrom, M., Sahlin, K. & Bangsbo, J. (2009). Four weeks of speed endurance training reduces energy expenditure during exercise and maintains muscle oxidative capacity despite a reduction in training volume. *J. Appl. Physiol.*, 106, 73-80.
- Impellizzeri, F. M., Marcora, S. M., Castagna, C., Reilly, T., Sassi, A. et al. (2006). Physiological and performance effects of generic versus specific aerobic training in soccer players. *Int. J. Sports Med.*, 27, 483-492.
- Laursen, P. B., Shing, C. M., Peake, J. M., Coombes, J. S. & Jenkins, D. G. (2002). Interval training program optimization in highly trained endurance cyclists. *Med. Sci. Sports Exerc.*, 34, 1801-1807.
- Laursen, P. B., Shing, C. M., Peake, J. M., Coombes, J. S. & Jenkins, D. G. (2005). Influence of high-intensity interval training on adaptations in well-trained cyclists. *J. Strength Cond. Res.*, 19, 527-533.
- McMillan, K., Helgerud, J., MacDonald, R. & Hoff, J. (2005). Physiological adaptations to soccer specific endurance training in professional youth soccer players. *Br. J. Sports Med.*, 39 (5), 273-277.
- McKenna, M., Heigenhauser, G., McKelvie, R. et al. (1997). Sprint training enhances ionic regulation during intense exercise in men. *J. Physiol.*, 515, 687-702.
- Mohr, M., Krstrup, P. & Bangsbo, J. (2003). Match performance of high-standard soccer players with special reference to development of fatigue. *J. Sports Sci.*, 18, 519-528.
- Nagler, P. & Gruber, S. (2004). *Fußballtraining Planung und Steuerung – Fit aus der Winterpause*. Aachen: Meyer und Meyer.
- Reilly, T. (1997). Energistics of high-intensity exercise (soccer) with particular reference to fatigue. *J. Sports Sci.*, 15, 257-263.
- Reilly, T. (2003). Motion analysis and physiological demands. In A. M. Williams & T. Reilly (eds.), *Science and Soccer* (2nd ed.; pp. 59-72). London: E and FN Spon.
- Reilly, T. (2005). An ergonomics model of the soccer training process. *J. Sport Sci.*, 13, 561-572.
- Stølen, T. O., Chamari, K., Castagna, C. & Wisloff, U. (2005). Physiology of soccer an update. *Sports Med.*, 35 (6), 501-536.
- Sperlich, B., Florian, E., Broich, H., Krüger, M., Zinner, C. et al. (2010). Vergleich von intensivem Intervalltraining vs. umfangsbetontem Ausdauertraining in der Vorbereitungsphase im U14-Fußball. *Sportmed. Sporttraumatol.*, 58 (4), 120-124.
- Stepto, N. K., Hawley, J. A., Dennis, S. C. & Hopkins, W. G. (1999). Effects of different interval-training programs on cycling time-trial performance. *Med. Sci. Sports Exerc.*, 31, 736-741.
- Sporis, G., Ruzic, L. & Leko, G. (2008). The anaerobic endurance of elite soccer players improved after a high-intensity training intervention in the 8-week conditioning program. *J. Strength Cond. Res.*, 22, 559-566.
- Tabata, I., Nishimura, M., Kouzaki, M., Hirai, Y. et al. (1996). Effects of moderate intensity-endurance and high intensity-intermittent training on anaerobic capacity and  $\text{VO}_{2\text{max}}$ . *Med. Sci. Sports Exerc.*, 28, 1327-1330.

Thorstensson, A., Sjödin, B. & Karlsson, J. (1975). Enzyme activities and muscle strength after "sprint training" in man. *Acta Physiol. Scand.*, 94 (3), 313-318.

Verheijen, R. (1997). *Fußballkondition*. Amsterdam: Wilco, Amersfoort. Deutsche Übersetzung von Winfried Schoofs (1999/2000).

Van Gool, D., Van Gerven, D. & Boutmans, J. (1988). The physiological load imposed on soccer players during real match-play. In T. Reilly, A. Lees, K. Davids et al. (eds.). *Science and Football* (pp. 51-59). London: Spon.

Wisloff, U., Helgerud, J. & Hoff, J. (1998). Strength and endurance of elite soccer players. *Med. Sci. Sports Exerc.*, 30, 462-467.

Weineck, A. & Weineck, J. (2005). *Leistungskurs Sport, Band II, Sportbiologische und trainingswissenschaftliche Grundlagen*. 2. erweiterte Auflage. Forchheim: Edition Zenk.

#### Korrespondenzadresse

Univ. Prof. Mag. Dr. Peter Hofmann, FACSM, Institute of Sports Science, Exercise Physiology & Training Research Group, University of Graz, Max-Mell-Allee 11, A-8010 Graz, Austria

E-Mail: peter.hofmann@uni-graz.at