

Alexander Törpel/Lutz Schega

AKTUELLE ENTWICKLUNGEN IM HÖHENTRAINING

Allgemeine und schwimmsportspezifische Erkenntnisse im narrativen Überblick

Literatur

- Balsalobre-Fernández, C., Tejero-González, C. M. & del Campo-Curiel, D. (2014). Hypoxic exposure as a means of increasing sporting performance: fact or fiction? *Revista Internacional de Medicina y Ciencias de la Actividad Física y el Deporte*, 14 (53), 183–198.
- Bärtsch, P., Dehnert, C., Friedmann-Bette, B. & Tadibi, V. (2008). Intermittent hypoxia at rest for improvement of athletic performance. *Scandinavian Journal of Medicine & Science in Sports*, 18, 50–56 (doi: 10.1111/j.1600-0838.2008.00832.x).
- Beidleman, B. A., Fulco, C. S., Staab, J. E., Andrew, S. P. & Muza, S. R. (2014). Cycling performance decrement is greater in hypobaric versus normobaric hypoxia. *Extreme Physiology & Medicine*, 3 (1), 1–9 (doi: 10.1186/2046-7648-3-8).
- Bejder, J. & Nordsborg, N. B. (2018). Specificity of "Live High-Train Low" Altitude Training on Exercise Performance. *Exercise and sport sciences reviews*, 46 (2), S. 129–136 (doi: 10.1249/JES.000000000000144).
- Berthelot, G., Sedeaud, A., Marck, A., Antero-Jacquemin, J., Schipman, J., Saulière, G., Marc, A., Desgorges, F.-D. & Toussaint, J.-F. (2015). Has Athletic Performance Reached its Peak? *Sports Medicine*, 45 (9), S. 1263–1271 (doi: 10.1007/s40279-015-0347-2).
- Bonetti, D. L. & Hopkins, W. G. (2009). Sea-level exercise performance following adaptation to hypoxia: a meta-analysis. *Sports medicine (Auckland, N.Z.)*, 39 (2), S. 107–127 (doi: 10.2165/00007256-200939020-00002).
- Bonne, T. C., Lundby, C., Jørgensen, S., Johansen, L., Mrgan, M., Bech, S. R., Sander, M., Papoti, M. & Nordsborg, N. B. (2014). "Live High-Train High" increases hemoglobin mass in Olympic swimmers. *European Journal of Applied Physiology*, 114 (7), 1439–1449 (doi: 10.1007/s00421-014-2863-4).
- Brocherie, F., Millet, G. P., Hauser, A., Steiner, T., Rysman, J., Wehrli, J. P. & Girard, O. (2015). "Live High-Train Low and High" Hypoxic Training Improves Team-Sport Performance. *Medicine & Science in Sports & Exercise*, 47 (10), 2140–2149 (doi: 10.1249/MSS.0000000000000630).
- Buhl, H. (2011). Die systematische Wirkung der Hypoxie auf den Organismus als Grundlage der Anwendung als Therapieform bei ausgewählten Krankheiten. In Kuno Hottenrott (Hrsg.), *Herzfreizugsvariabilität. Gesundheitsförderung, Trainingssteuerung, Biofeedback: internationales Symposium vom 29. und 30. Oktober 2010 in Halle (Saale) (Schriften der Deutschen Vereinigung für Sportwissenschaft, 214, S. 11–23)*. Hamburg: Czwalina Verlag
- Burtscher, M. (2010). Auswirkungen akuter Höhenexposition: Welche Höhe ist gesundheitlich verträglich? *Wiener Medizinische Wochenschrift*, 160 (13–14), 362–371 (doi: 10.1007/s10354-010-0742-0).
- Burtscher, M., Gatterer, H., Faulhaber, M., Gerstgrasser, W. & Schenk, K. (2010). Effects of intermittent hypoxia on running economy. *International journal of sports medicine*, 31 (9), 644–650 (doi: 10.1055/s-0030-1255067).
- Burtscher, M., Pachinger, O., Ehrenbourg, I., Mitterbauer, G., Faulhaber, M., Pühringer, R. & Tkatchouk, E. (2004). Intermittent hypoxia increases exercise tolerance in elderly men with and without coronary artery disease. *International journal of cardiology*, 96 (2), S. 247–254 (doi: 10.1016/j.ijcard.2003.07.021).
- Chapman, R. F. (2013). The individual response to training and competition at altitude. *British journal of sports medicine*, 47 (Suppl. 1), i40–44 (doi: 10.1136/bjsports-2013-092837).
- Chapman, R. F., Laymon Stickford, A. S., Lundby, C. & Levine, B. D. (2014). Timing of return from altitude training for optimal sea level performance. *Journal of Applied Physiology*, 116 (7), 837–843 (doi: 10.1152/jappphysiol.00663.2013).
- Chapman, R. F., Stray-Gundersen, J. & Levine, B. D. (1998). Individual variation in response to altitude training. *Journal of Applied Physiology*, 85 (4), 1448–1456.
- Conkin, J. & Wessel, J. H. (2008). Critique of the Equivalent Air Altitude Model. *Aviation, space, and environmental medicine*, 79 (10), 975–982 (doi: 10.3357/ASEM.2331.2008).
- Czuba, M., Wilk, R., Karpiński, J., Chalimoniuk, M., Zajac, A. & Langfort, J. (2017). Intermittent hypoxic training improves anaerobic performance in competitive swimmers when implemented into a direct competition mesocycle. *PLoS ONE*, 12 (8), e0180380 (doi: 10.1371/journal.pone.0180380).
- Deb, S. K., Brown, D. R., Gough, L. A., McLellan, C. P., Swinton, P. A., Andy Sparks, S. & McNaughton, L. R. (2018). Quantifying the effects of acute hypoxic exposure on exercise performance and capacity: A systematic review and meta-regression. *European Journal of Sport Science*, 18 (2), 243–256 (doi: 10.1080/17461391.2017.1410233).
- Flaherty, G., O'Connor, R. & Johnston, N. (2016). Altitude training for elite endurance athletes: A review for the travel medicine practitioner. *Travel medicine and infectious disease*, 14 (3), S. 200–211 (doi: 10.1016/j.tmaid.2016.03.015).
- Frese, F. & Friedmann-Bette, B. (2010). Effects of repetitive training at low altitude on erythropoiesis in 400 and 800 m runners. *International journal of sports medicine*, 31 (6), 382–388 (doi: 10.1055/s-0030-1248328).
- Friedmann, B., Frese, F., Menold, E., Kauper, F., Jost, J. & Bärtsch, P. (2005). Individual variation in the erythropoietic response to altitude training in elite junior swimmers. *British journal of sports medicine*, 39 (3), 148–153 (doi: 10.1136/bjism.2003.011387).
- Fulco, C. S., Rock, P. B. & Cymerman, A. (1998). Maximal and submaximal exercise performance at altitude. *Aviation, space, and environmental medicine*, 69 (8), 793–801.
- Garvican, L., Martin, D., Quod, M., Stephens, B., Sassi, A. & Gore, C. (2012). Time course of the hemoglobin mass response to natural altitude training in elite endurance cyclists. *Scandinavian Journal of Medicine & Science in Sports*, 22 (1), 95–103 (doi: 10.1111/j.1600-0838.2010.01145.x).
- Garvican-Lewis, L. A., Govus, A. D., Peeling, P., Abbiss, C. R. & Gore, C. J. (2016). Iron Supplementation and Altitude. Decision Making Using a Regression Tree. *Journal of sports science & medicine*, 15 (1), 204–205.
- Girard, O., Amann, M., Aughey, R., Billaut, F., Bishop, D. J., Bourdon, P., Buchheit, M., Chapman, R., D'Hooghe, M., Garvican-Lewis, L. A., Gore, C. J., Millet, G. P., Roach, G. D., Sargent, C., Saunders, P. U., Schmidt, W. & Schumacher, Y. O. (2013). Position statement—altitude training for improving team-sport players' performance: current knowledge and unresolved issues. *British journal of sports medicine*, 47 (Suppl. 1), i8–i16 (doi: 10.1136/bjsports-2013-093109).
- Girard, O., Brocherie, F. & Millet, G. P. (2017). Effects of Altitude/Hypoxia on Single- and Multiple-Sprint Performance: A Comprehensive Review. *Sports medicine (Auckland, N.Z.)*, 47 (10), 1931–1949 (doi: 10.1007/s40279-017-0733-z).
- Girard, O., Koehle, M. S., MacInnis, M. J., Guenette, J. A., Verges, S., Rupp, T., Jubeau, M., Perrey, S., Millet, G. Y., Chapman, R. F., Levine, B. D., Conkin, J., Wessel, J. H., Nespoulet, H., Wuyam, B., Tamisier, R., Levy, P., Casey, D. P., Taylor, B. J., Snyder, E. M., Johnson, B. D., Laymon, A. S., Stickford, J. L., Weavil, J. C., Loeppky, J. A., Pun, M., Schommer, K., Bartsch, P., Vagula, M. C. & Nelatury, C. F. (2012). Comments on Point: Counterpoint: Hypobaric hypoxia induces/does not induce different responses from normobaric hypoxia. *Journal of Applied Physiology*, 112 (10), 1788–1794 (doi: 10.1152/jappphysiol.00356.2012).
- Gore, C. J., Clark, S. A. & Saunders, P. U. (2007). Non-hematological mechanisms of improved sea-level performance after hypoxic exposure. *Medicine and science in sports and exercise*, 39 (9), 1600–1609 (doi: 10.1249/mss.0b013e3180de49d3).
- Gore, C. J., Sharpe, K., Garvican-Lewis, L. A., Saunders, P. U., Humberstone, C. E., Robertson, E. Y., Wachsmuth, N. B., Clark, S. A., McLean, B. D., Friedmann-Bette, B., Neya, M., Pottgiesser, T., Schumacher, Y. O. & Schmidt, W. F. (2013). Altitude training and haemoglobin mass from the optimised carbon monoxide rebreathing method determined by a meta-analysis. *British journal of sports medicine*, 47 (Suppl. 1), i31–i39 (doi: 10.1136/bjsports-2013-092840).
- Gough, C. E., Saunders, P. U., Fowlie, J., Savage, B., Pyne, D. B., Anson, J. M., Wachsmuth, N., Prommer, N. & Gore, C. J. (2012). Influence of altitude training modality on performance and total haemoglobin mass in elite swimmers. *European Journal of Applied Physiology*, 112 (9), 3275–3285 (doi: 10.1007/s00421-011-2291-7).
- Govus, A. D., Garvican-Lewis, L. A., Abbiss, C. R., Peeling, P. & Gore, C. J. (2015). Pre-Altitude Serum Ferritin Levels and Daily Oral Iron Supplement Dose

- Mediate Iron Parameter and Hemoglobin Mass Responses to Altitude Exposure. *PLoS ONE*, 10 (8), e0135120 (doi: 10.1371/journal.pone.0135120).
- Green, H., MacDougall, J., Tarnopolsky, M. & Melissa, N. L. (1999). Downregulation of Na⁺-K⁺-ATPase pumps in skeletal muscle with training in normobaric hypoxia. *Journal of Applied Physiology*, 86 (5), 1745–1748.
- Harshman, S. W., Geier, B. A., Fan, M., Rinehardt, S., Watts, B. S., Drummond, L. A., Preti, G., Phillips, J. B., Ott, D. K. & Grigsby, C. C. (2015). The identification of hypoxia biomarkers from exhaled breath under normobaric conditions. *Journal of Breath Research*, 9 (4), 47103 (doi: 10.1088/1752-7155/9/4/047103).
- Hauser, A., Schmitt, L., Troesch, S., Saugy, J. J., Cejuela-Anta, R., Faiss, R., Robinson, N., Wehrin, J. P. & Millet, G. P. (2016). Similar Hemoglobin Mass Response in Hypobaric and Normobaric Hypoxia in Athletes. *Medicine and Science in Sports and Exercise*, 48 (4), 734–741 (doi: 10.1249/MSS.0000000000000808).
- Hecksteden, A., Faude, O., Meyer, T. & Donath, L. (2018). How to Construct, Conduct and Analyze an Exercise Training Study? *Frontiers in physiology*, 9, 1007 (doi: 10.3389/fphys.2018.01007).
- Heinicke, K., Hofer, T., Wenger, R.H. & Gassmann, M. (2002). Die zelluläre Antwort auf Sauerstoffmangel. The cellular response to hypoxia. *Deutsche Zeitschrift für Sportmedizin*, 53 (10), 270–276.
- Holliss, B. A., Burden, R. J., Jones, A. M. & Pedlar, C. R. (2014). Eight weeks of intermittent hypoxic training improves submaximal physiological variables in highly trained runners. *Journal of Strength and Conditioning Research*, 28 (8), 2195–2203 (doi: 10.1519/JSC.0000000000000406).
- Hoppeler, H. & Vogt, M. (2001). Muscle tissue adaptations to hypoxia. *The Journal of experimental biology*, 204 (Pt 18), 3133–3139.
- Jedlickova, K., Stockton, D. W., Chen, H., Stray-Gundersen, J., Witkowski, S., Ri-Li, G., Jelinek, J., Levine, B. D. & Prchal, J. T. (2003). Search for genetic determinants of individual variability of the erythropoietin response to high altitude. *Blood Cells, Molecules, and Diseases*, 31 (2), 175–182 (doi: 10.1016/S1079-9796(03)00153-0).
- Keslacy, S. (2006). Comments on Point-Counterpoint "Positive effects of intermittent hypoxia (live high:train low) on exercise performance are/are not mediated primarily by augmented red cell volume". *Journal of Applied Physiology*, 100 (1), 363 (doi: 10.1152/jappphysiol.01156.2005).
- Levine, B. D. (2002). Intermittent hypoxic training: fact and fancy. *High altitude medicine & biology*, 3 (2), 177–193 (doi: 10.1089/15270290260131911).
- Levine, B. D. & Stray-Gundersen, J. (1992). A practical approach to altitude training: where to live and train for optimal performance enhancement. *International journal of sports medicine*, 13 (Suppl. 1), 209–212 (doi: 10.1055/s-2007-1024642).
- Levine, B. D. & Stray-Gundersen, J. (1997). "Living high-training low": effect of moderate-altitude acclimatization with low-altitude training on performance. *Journal of Applied Physiology*, 83 (1), 102–112.
- Levine, B. D. & Stray-Gundersen, J. (2005). Point: Positive effects of intermittent hypoxia (live high:train low) on exercise performance are mediated primarily by augmented red cell volume. *Journal of Applied Physiology*, 99 (5), 2053–2055 (doi: 10.1152/jappphysiol.00877.2005).
- McLean, B. D., Buttifant, D., Gore, C. J., White, K. & Kemp, J. (2013a). Year-to-year variability in haemoglobin mass response to two altitude training camps. *British journal of sports medicine*, 47 (Suppl. 1), i51–58 (doi: 10.1136/bjsports-2013-092744).
- McLean, B. D., Buttifant, D., Gore, C. J., White, K., Liess, C. & Kemp, J. (2013b). Physiological and performance responses to a preseason altitude-training camp in elite team-sport athletes. *International journal of sports physiology and performance*, 8 (4), 391–399.
- Millet, G. P., Faiss, R., Brocherie, F. & Girard, O. (2013). Hypoxic training and team sports: a challenge to traditional methods? *British journal of sports medicine*, 47 (Suppl. 1), i6–7 (doi: 10.1136/bjsports-2013-092793).
- Millet, G. P., Faiss, R. & Pialoux, V. (2012). Point: Counterpoint: Hypobaric hypoxia induces/does not induce different responses from normobaric hypoxia. *Journal of Applied Physiology*, 112 (10), 1783–1784 (doi: 10.1152/jappphysiol.00067.2012).
- Millet, G. P., Roels, B., Schmitt, L., Woorons, X. & Richalet, J. P. (2010). Combining hypoxic methods for peak performance. *Sports Medicine*, 40 (1), 1–25 (doi: 10.2165/11317920-000000000-00000).
- Moon, H.-W., Sunoo, S., Park, H.-Y., Lee, D.-J. & Nam, S.-S. (2016). Effects of various acute hypoxic conditions on metabolic parameters and cardiac function during exercise and recovery. *SpringerPlus*, 5 (1), 1294 (doi: 10.1186/s40064-016-2952-4).
- Mounier, R. & Brugniaux, J. V. (2012). Counterpoint: Hypobaric hypoxia does not induce different responses from normobaric hypoxia. *Journal of Applied Physiology*, 112 (10), 1784–1786 (doi: 10.1152/jappphysiol.00067.2012a).
- Mourou, L. & Millet, G. P. (2019). Is Maximal Heart Rate Decrease Similar Between Normobaric Versus Hypobaric Hypoxia in Trained and Untrained Subjects? *High Alt Med Biol.*, 20 (1), 94–98 (doi: 10.1089/ham.2018.0104).
- Netzer, N. C., Rausch, L., Eliasson, A. H., Gatterer, H., Friess, M., Burtcher, M. & Pramsohler, S. (2017). SpO₂ and Heart Rate During a Real Hike at Altitude Are Significantly Different than at Its Simulation in Normobaric Hypoxia. *Frontiers in physiology*, 8, 81 (doi: 10.3389/fphys.2017.00081).
- Nieß, A. (2010). Sport in der Höhe und unter Hypoxiebedingungen. Klassisches Höhentraining. In H.-H. Dickhuth, F. Mayer, K. Röcker & A. Berg (Hrsg.): *Sportmedizin für Ärzte. Lehrbuch auf der Grundlage des Weiterbildungssystems der Deutschen Gesellschaft für Sportmedizin und Prävention (DGSP)*. 2. Aufl. Köln: Deutscher Ärzte Verlag.
- Noakes, T. D. (2005). Comments on Point:Counterpoint "Positive effects of intermittent hypoxia (live high:train low) on exercise performance are/are not mediated primarily by augmented red cell volume". *Journal of Applied Physiology*, 99 (6), 2453 (doi: 10.1152/jappphysiol.01157.2005).
- Ofner, M., Wonisch, M., Frei, M., Tschakert, G., Domej, W., Kröpfel, J. M. & Hofmann, P. (2014). Influence of Acute Normobaric Hypoxia on Physiological Variables and Lactate Turn Point Determination in Trained Men. *Journal of sports science & medicine*, 13 (4), 774–781.
- Park, H.-Y. & Lim, K. (2017). Effects of Hypoxic Training versus Normoxic Training on Exercise Performance in Competitive Swimmers. *Journal of sports science & medicine*, 16 (4), 480–488.
- Park, H.-Y., Shin, C. & Lim, K. (2018). Intermittent hypoxic training for 6 weeks in 3000 m hypobaric hypoxia conditions enhances exercise economy and aerobic exercise performance in moderately trained swimmers. *Biology of Sport*, 35 (1), 49–56 (doi: 10.5114/biolsport.2018.70751).
- Paula, P. de & Niebauer, J. (2012). Effects of high altitude training on exercise capacity. Fact or myth. *Sleep & breathing = Schlaf & Atmung*, 16 (1), 233–239 (doi: 10.1007/s11325-010-0445-1).
- Pedlar, C. R., Whyte, G. P. & Godfrey, R. J. (2008). Pre-acclimation to exercise in normobaric hypoxia. *European Journal of Sport Science*, 8 (1), 15–21 (doi: 10.1080/17461390701871932).
- Pinilla, V. & Cecilia, O. (2014). Exercise and Training at Altitudes: Physiological Effects and Protocols. *Revista Ciencias de la Salud*, 12 (1), 115–130 (doi: 10.12804/revsalud12.1.2014.07).
- Płoszczyca, K., Langfort, J. & Czuba, M. (2018). The Effects of Altitude Training on Erythropoietic Response and Hematological Variables in Adult Athletes. A Narrative Review. *Frontiers in physiology*, 9, 375 (doi: 10.3389/fphys.2018.00375).
- Ponsot, E., Dufour, S. P., Zoll, J., Doutrelau, S., N'Guesan, B., Geny, B., Hoppeler, H., Lampert, E., Mettauer, B., Ventura-Clapier, R. & Richard, R. (2006). Exercise training in normobaric hypoxia in endurance runners. II. Improvement of mitochondrial properties in skeletal muscle. *Journal of Applied Physiology*, 100 (4), 1249–1257 (doi: 10.1152/jappphysiol.00361.2005).
- Rebuttal from Millet, Faiss, and Pialoux (2012). *Journal of applied physiology (Bethesda, Md. : 1985)*, 112 (10), 1786–1787.
- Richard, N. A. & Koehle, M. S. (2012). Differences in Cardio-Ventilatory Responses to Hypobaric and Normobaric Hypoxia: A Review. *Aviation, space, and environmental medicine*, 83 (7), 677–684 (doi: 10.3357/ASEM.3182.2012).
- Robach, P., Schmitt, L., Brugniaux, J. V., Roels, B., Millet, G., Hellard, P., Nicolet, G., Duvallet, A., Fouillot, J.-P., Moutereau, S., Lasne, F., Pialoux, V., Olsen, N. V. & Richalet, J.-P. (2006). Living high-training low: effect on erythropoiesis and aerobic performance in highly-trained swimmers. *European Journal of Applied Physiology*, 96 (4), 423–433 (doi: 10.1007/s00421-005-0089-1).
- Robertson, E. Y., Aughey, R. J., Anson, J. M., Hopkins, W. G. & Pyne, D. B. (2010a). Effects of Simulated and Real Altitude Exposure in Elite Swimmers. *Journal of Strength and Conditioning Research*, 24 (2), 487–493 (doi: 10.1519/JSC.0b013e3181c06d56).
- Robertson, E. Y., Saunders, P. U., Pyne, D. B., Aughey, R. J., Anson, J. M. & Gore, C. J. (2010b). Reproducibility of Performance Changes to Simulated Live High/Train Low Altitude. *Medicine & Science in Sports & Exercise*, 42 (2), 394–401 (doi: 10.1249/MSS.0b013e3181b34b57).
- Robertson, E. Y., Saunders, P. U., Pyne, D. B., Gore, C. J. & Anson, J. M. (2010c). Effectiveness of intermittent training in hypoxia combined with live high/train low. *European journal of applied physiology*, 110 (2), 379–387 (doi: 10.1007/s00421-010-1516-5).
- Rodriguez, F. A., Truijens, M. J., Townsend, N. E., Stray-Gundersen, J., Gore, C. J. & Levine, B. D. (2007). Performance of runners and swimmers after four weeks of intermittent hypobaric hypoxic exposure plus sea level training. *Journal of Applied Physiology*, 103 (5), 1523–1535 (doi: 10.1152/jappphysiol.01320.2006).
- Rodríguez, F. A., Iglesias, X., Feriche, B., Caldersón-Soto, C., Chaverri, D., Wachsmuth, N. B., Schmidt, W. & Levine, B. D. (2015). Altitude Training in Elite Swimmers for Sea Level Performance (Altitude Project). *Medicine and Science in Sports and Exercise*, 47 (9), 1965–1978 (doi: 10.1249/MSS.0000000000000626).
- Rodríguez, F. A., Murio, J. & Ventura, J. L. (2003). Effects of intermittent hypobaric hypoxia and altitude training on physiological and performance parameters in swimmers. *Medicine & Science in Sports & Exercise*, 35 (5), 115.
- Roels, B., Hellard, P., Schmitt, L., Robach, P., Richalet, J. P. & Millet, G. P. (2006). Is it more effective for highly trained swimmers to live and train at 1200 m than at 1850 m in terms of performance and haematological benefits? *British Journal of Sports Medicine*, 40 (2), e4 (doi: 10.1136/bjism.2004.017103).
- Ross, R., Goodpaster, B. H., Koch, L. G., Sarzynski, M. A., Kohrt, W. M., Johannsen, N. M., Skinner, J. S., Castro, A., Irving, B. A., Noland, R. C., Sparks, L. M., Spielmann, G., Day, A. G., Pitsch, W., Hopkins, W. G. & Boucharde, C. (2019). Precision exercise medicine: understanding exercise response variability. *British journal of sports medicine* (doi: 10.1136/bjsports-2018-100328).

- Saugy, J. J., Schmitt, L., Cejuela, R., Faiss, R., Hauser, A., Wehrlin, J. P., Rudaz, B., Delessert, A., Robinson, N. & Millet, G. P. (2014). Comparison of "Live High-Train Low" in normobaric versus hypobaric hypoxia. *PLoS ONE*, 9 (12), e114418 (doi: 10.1371/journal.pone.0114418).
- Saugy, J. J., Schmitt, L., Hauser, A., Constantin, G., Cejuela, R., Faiss, R., Wehrlin, J. P., Rosset, J., Robinson, N. & Millet, G. P. (2016). Same Performance Changes after Live High-Train Low in Normobaric vs. Hypobaric Hypoxia. *Frontiers in Physiology*, 7, 107 (doi: 10.3389/fphys.2016.00138).
- Saunders, P. U., Garvican-Lewis, L. A., Schmidt, W. F. & Gore, C. J. (2013). Relationship between changes in haemoglobin mass and maximal oxygen uptake after hypoxic exposure. *British Journal of Sports Medicine*, 47 (1), i26-i30 (doi: 10.1136/bjsports-2013-092841).
- Saunders, P. U., Pyne, D. B. & Gore, C. J. (2009). Endurance training at altitude. *High altitude medicine & biology*, 10 (2), 135-148 (doi: 10.1089/ham.2008.1092).
- Schmidt, W. & Prommer, N. (2010). Impact of alterations in total hemoglobin mass on VO_2max . *Exercise and sport sciences reviews*, 38 (2), 68-75 (doi: 10.1097/JES.0b013e3181d4957a).
- Schmitt, L., Millet, G., Robach, P., Nicolet, G., Brugniaux, J. V., Fouillat, J.-P. & Richalet, J.-P. (2006). Influence of "living high-training low" on aerobic performance and economy of work in elite athletes. *European Journal of Applied Physiology*, 97 (5), 627-636 (doi: 10.1007/s00421-006-0228-3).
- Schmitt, L., Willis, S. J., Coulmy, N. & Millet, G. P. (2018). Effects of Different Training Intensity Distributions Between Elite Cross-Country Skiers and Nordic-Combined Athletes During Live High-Train Low. *Frontiers in physiology*, 9, 932 (doi: 10.3389/fphys.2018.00932).
- Schuler, B., Lundby, C. & Gassmann, M. (2005). HIF-1 and the adaptation of man to high altitude. *Schweizerische Zeitschrift für Sportmedizin und Sporttraumatologie*, 53 (2), S. 82-87.
- Schumacher, Y. O. & Pottgießer, T. (2010). Höhentaining - Aktueller Stand und neue Tendenzen. *Leistungssport*, 40 (6), 8-12.
- Siewierski, M., Stomiński, P., Biatacki, R. & Adamczyk, J. (2012). Athletic performance of swimmers after altitude training (2.300m above sea level) in view of their blood morphology changes. *Biology of Sport*, 29 (2), 115-120 (doi: 10.5604/20831862.990749).
- Sinex, J. A. & Chapman, R. F. (2015). Hypoxic training methods for improving endurance exercise performance. *Journal of Sport and Health Science*, 4 (4), 325-332 (doi: 10.1016/j.jshs.2015.07.005).
- Stanula, A., Maszczyk, A., Roczniok, R., Pietraszewski, P., Ostrowski, A., Zajac, A. & Strzata, M. (2012). The development and prediction of athletic performance in freestyle swimming. *Journal of Human Kinetics*, 32, 97-107 (doi: 10.2478/v10078-012-0027-3).
- Strzata, M., Ostrowski, A. & Szyguta, Z. (2011). Altitude Training and its Influence on Physical Endurance in Swimmers. *Journal of Human Kinetics*, 28 (-1). DOI: 10.2478/v10078-011-0026-9).
- Terrados, N., Jansson, E., Sylvén, C. & Kaijser, L. (1990). Is hypoxia a stimulus for synthesis of oxidative enzymes and myoglobin? *Journal of Applied Physiology*, 68 (6), 2369-2372.
- Törpel, A., Peter, B., Hamacher, D. & Schega, L. (2019). Dose-response relationship of intermittent normobaric hypoxia to stimulate erythropoietin in the context of health promotion in young and old people. *European Journal of Applied Physiology* (doi: 10.1007/s00421-019-04096-8).
- Trincat, L., Woorons, X. & Millet, G. P. (2017). Repeated-Sprint Training in Hypoxia Induced by Voluntary Hypoventilation in Swimming. *International journal of sports physiology and performance*, 12 (3), 329-335 (doi: 10.1123/ijspp.2015-0674).
- Truijens, M. J., Rodríguez, F. A., Townsend, N. E., Stray-Gundersen, J., Gore, C. J. & Levine, B. D. (2008). The effect of intermittent hypobaric hypoxic exposure and sea level training on submaximal economy in well-trained swimmers and runners. *Journal of Applied Physiology*, 104 (2), 328-337 (doi: 10.1152/jappphysiol.01324.2006).
- Truijens, M. J., Toussaint, H. M., Dow, J. & Levine, B. D. (2003). Effect of high-intensity hypoxic training on sea-level swimming performances. *Journal of Applied Physiology*, 94 (2), 733-743 (doi: 10.1152/jappphysiol.00079.2002).
- Turner, G., Gibson, O. R., Watt, P. W., Pringle, J. S. M., Richardson, A. J. & Maxwell, N. S. (2017). The time course of endogenous erythropoietin, IL-6, and TNF in response to acute hypoxic exposures. *Scandinavian journal of medicine & science in sports*, 27 (7), 714-723 (doi: 10.1111/sms.12700).
- Vellers, H. L., Kleeberger, S. R. & Lightfoot, J. T. (2018). Inter-individual variation in adaptations to endurance and resistance exercise training: genetic approaches towards understanding a complex phenotype. *Mammalian genome: official journal of the International Mammalian Genome Society*, 29 (1-2), 48-62 (doi: 10.1007/s00335-017-9732-5).
- Vogt, M. & Hoppeler, H. (2010). Is hypoxia training good for muscles and exercise performance? *Progress in cardiovascular diseases*, 52 (6), 525-533 (doi: 10.1016/j.pcad.2010.02.013).
- Vogt, M., Montagne, J. & Hoppeler, H. (2005). Live low - train high: muskuläre Aspekte und Leistungsfähigkeit. *Schweizerische Zeitschrift für Sportmedizin und Sporttraumatologie*, 53 (2), S. 76-81.
- Vogt, M., Puntschart, A., Geiser, J., Zuleger, C., Billeter, R. & Hoppeler, H. (2001). Molecular adaptations in human skeletal muscle to endurance training under simulated hypoxic conditions. *Journal of Applied Physiology*, 91 (1), S. 173-182.
- Wachsmuth, N. B., Voelzke, C., Hoffmann, J., Saunders, P. U. & Schmidt, W. F. (2013a). Leistungsentwicklung nach einem Höhentrainingslager - eine Pilotstudie. *Deutsche Zeitschrift für Sportmedizin*, 64 (11), 339-342 (doi: 10.5960/dzsm.2013.100).
- Wachsmuth, N. B., Voelzke, C., Prommer, N., Schmidt-Trucksäss, A., Frese, F., Spahl, O., Eastwood, A., Stray-Gundersen, J. & Schmidt, W. (2013b). The effects of classic altitude training on hemoglobin mass in swimmers. *European Journal of Applied Physiology*, 113 (5), 1199-1211 (doi: 10.1007/s00421-012-2536-0).
- Wehrlin, J. P. (2005). «Live high - train low»: Ein erfolgreiches Höhentrainingsparadigma zur Leistungssteigerung bei Eliteausdauerathleten? *Schweizerische Zeitschrift für Sportmedizin und Sporttraumatologie*, 53 (2), S. 68-75.
- Wehrlin, J. P. & Hallén, J. (2006). Linear decrease in VO_2max and performance with increasing altitude in endurance athletes. *European Journal of Applied Physiology*, 96 (4), 404-412 (doi: 10.1007/s00421-005-0081-9).
- Wilber, R. L. (2007). Application of altitude/hypoxic training by elite athletes. *Medicine & Science in Sports & Exercise*, 39 (9), 1610-1624 (doi: 10.1249/mss.0b013e3180de49e6).
- Wilber, R. L. (2011). Application of altitude/hypoxic training by elite athletes. *Journal of Human Sport and Exercise*, 6 (2), 271-286 (doi: 10.4100/jhse.2011.62.07).
- Wilber, R. L., Stray-Gundersen, J. & Levine, B. D. (2007). Effect of hypoxic "dose" on physiological responses and sea-level performance. *Medicine & Science in Sports & Exercise*, 39 (9), 1590-1599 (doi: 10.1249/mss.0b013e3180de49bd).
- Witkowski, S., Chen, H., Stray-Gundersen, J., Ge, R. L., Alfrey, C., Prchal, J. T. & Levine, B. D. (2002). Genetic Marker for the Erythropoietic Response to Altitude. *Medicine and science in sports and exercise*, 34 (5), 246 (doi: 10.1097/00005768-200205001-01375).
- Woorons, X., Mollard, P., Pichon, A., Lamberto, C., Duvallat, A. & Richalet, J.-P. (2007). Moderate exercise in hypoxia induces a greater arterial desaturation in trained than untrained men. *Scandinavian Journal of Medicine and Science in Sports*, 17 (4), 431-436 (doi: 10.1111/j.1600-0838.2006.00577.x).
- Zhong, N., Zhang, Y., Zhu, H. F. & Zhou, Z. N. (2000). Intermittent hypoxia exposure prevents mtDNA deletion and mitochondrial structure damage produced by ischemia/reperfusion injury. *Sheng li xue bao [Acta physiologica Sinica]*, 52 (5), 375-380.
- Zoretic, D., Grcic-Zubcevic, N. & Zubcic, K. (2014). The effects of hypercapnic-hypoxic training program on hemoglobin concentration and maximum oxygen uptake of elite swimmers. *Kinesiology*, 46, 40-45.
- Zoretic, D. & Kuterovac, P. (2012). Effect of 21 days of "Living High - Training High" training period on sea-level swimming performance and haematological blood results in competitive swimmers. *Gazzetta Medica Italiana - Archivio Per Le Scienze Mediche* (171), 1-9.

Korrespondenzadresse

Dr. phil. Alexander Törpel, Wissenschaftlicher Mitarbeiter an der Otto-von-Guericke-Universität Magdeburg, Institut III – Sportwissenschaft, Lehrstuhl Gesundheit und körperliche Aktivität, Zschokkestraße 32, 39104 Magdeburg
E-Mail: alexandertoerpel@ovgu.de