

Steffen Held/Michael Behringer/Lars Donath

BLUTFLUSSRESTRIKTION ERHÖHT DIE VO₂MAX BEI GUT TRAINIERTEN RUDERERN

Eine randomisiert-kontrollierte Studie beim Rudern mit geringer Intensität

Literatur

- Abe, T., Fujita, S., Nakajima, T. et al. (2010). Effects of low-intensity cycle training with restricted leg blood flow on thigh muscle volume and VO₂max in young men. *J. Sports Sci. Med.*, 9 (3), 452-458.
- Abe, T., Sakamaki, M., Fujita, S. et al. (2010). Effects of low-intensity walk training with restricted leg blood flow on muscle strength and aerobic capacity in older adults. *J. Geriatr. Phys. Ther.*, 33 (1), 34-40.
- Allison, R. C. & Lawrence Bedsole, D. (2003). The other medical causes of rhabdomyolysis. *Am. J. Med. Sci.*, 326 (2), 79-88.
- Behringer, M., Behlau, D., Montag, J. C. K. et al. (2017). Low-intensity sprint training with blood flow restriction improves 100-m dash. *J. Strength Cond. Res.*, 31 (9), 2462-2472.
- Bennett, H. & Slattery, F. (2018). Effects of blood flow restriction training on aerobic capacity and performance: a systematic review. *J. Strength Cond. Res.*, 33 (2), 572-583.
- Bourgeois, J., Steyaert, A. & Boone, J. (2014). Physiological and anthropometric progression in an international oarsman: a 15-year case study. *Int. J. Sports Physiol. Perform.*, 9 (4), 723-726.
- Cohen, J. (1988). *Statistical Power Analysis for the Behavioral Sciences*. Hillsdale: Lawrence Erlbaum Associates.
- Dickhuth, H.-H., Huonker, M., Münzel, T. et al. (1991). *Advances in Ergometry*. https://scholar.google.com/scholar_lookup?title=Advances+in+Ergometry&author=H.+H.+Dickhuth&author=M.+Huonker&author=T.+Münzel&author=H.+Drexler&author=A.+Berg&publication_year=1991&
- Guellich, A. & Schmidtbleicher, D. (1999). Struktur der Kraftfähigkeiten und ihrer Trainingsmethoden. *Dtsch. Z. Sportmed.*, 50 (7/8), <https://www.bisp-surf.de/Record/PU199912403468>.
- Harris, D. & Atkinson, G. (2015). Ethical standards in sport and exercise science research: 2016 update. *Int. J. Sports Med.*, 36 (14), 1121-1124.
- Hawley, J. A. (2009). Molecular responses to strength and endurance training: Are they incompatible? *Appl. Physiol. Nutr. Metab.*, 34 (3), 355-361.
- Hughes, L., Paton, B., Rosenblatt, B., Gissane, C., & Patterson, S. D. (2017). Blood flow restriction training in clinical musculoskeletal rehabilitation: a systematic review and meta-analysis. *British Journal of Sports Medicine*, 51 (13), 1003-1011.
- Ingham, S., Whyte, G., Jones, K. & Nevill, A. (2002). Determinants of 2,000 m rowing ergometer performance in elite rowers. *Eur. J. Appl. Physiol.*, 88 (3), 243-246.
- Iversen, E., & Røstad, V. (2010). Low-load ischemic exercise-induced rhabdomyolysis. *Clinical Journal of Sport Medicine: Official Journal of the Canadian Academy of Sport Medicine*, 20 (3), 218-219.
- Izquierdo-Gabarrén, M., González de Txabarri Expósito, R., García-Pallarés, J. et al. (2009). Concurrent endurance and strength training not to failure optimizes performance gains. *Med. Sci. Sport Exerc.*, 42 (6): 1 (doi:10.1249/MSS.0b013e3181c67eec).
- Kraemer, W. J., Fry, A. C., Maud, P. & Foster, C. (1995). *Strength Testing: Development and Evaluation of Methodology*. Champaign (IL): Human Kinetics.
- Lixandrão, M. E., Ugrinowitsch, C., Berton, R. et al. (2018). Magnitude of muscle strength and mass adaptations between high-load resistance training versus low-load resistance training associated with blood-flow restriction: a systematic review and meta-analysis. *Sport Med.*, 48 (2), 361-378.
- Loenneke, J. P., Wilson, J. M., Marín, P. J. et al. (2012). Low intensity blood flow restriction training: a meta-analysis. *Eur. J. Appl. Physiol.*, 112 (5), 1849-1859.
- Loenneke, J. P., Wilson, J. M., Wilson, G. J. et al. (2011). Potential safety issues with blood flow restriction training. *Scand. J. Med. Sci. Sports*, 21 (4), 510-518.
- Macfarlane, D. J. & Wong, P. (2012). Validity, reliability and stability of the portable Cortex Metamax 3B gas analysis system. *Eur. J. Appl. Physiol.*, 112 (7), 2539-2547.
- Matsumoto, S. T. S. A., Suzuki, Y., Azuma, K., & Matsumoto, H. (2016). Rhabdomyolysis after performing blood flow restriction training: A case report. *J. Strength Cond. Res.*, 30 (7), 2064-2068.
- McBride, J. M., Triplett-McBride, T., Davie, A. & Newton, R. U. (2002). The effect of heavy- vs. light-load jump squats on the development of strength, power, and speed. *J. Strength Cond. Res.*, 16 (1), 75-82.
- Midgley, A. W., McNaughton, L. R., Polman, R. & Marchant, D. (2007). Criteria for determination of maximal oxygen uptake. *Sport Med.*, 37 (12), 1019-1028.
- Nakajima, T., Kurano, M., Lida, H., Takano, H., Oonuma, H., Morita, T., ... KAATSU Training Group, K. T. (2006). Use and safety of KAATSU training: Results of a national survey. *International Journal of KAATSU Training Research*, 2 (1), 5-13.
- Park, S., Kim, J. K., Choi, H. M. et al. (2010). Increase in maximal oxygen uptake following 2-week walk training with blood flow occlusion in athletes. *Eur. J. Appl. Physiol.*, 109 (4), 591-600.
- Patterson, S. D., Hughes, L., Warmington, S. et al. (2019). Blood flow restriction exercise position stand: considerations of methodology, application, and safety. *Front Physiol.*, 10: 533 (doi:10.3389/fphys.2019.00533).
- Pripstein, L. P., Rhodes, E. C., McKenzie, D. C. & Coutts, K. D. (1999). Aerobic and anaerobic energy during a 2-km race simulation in female rowers. *Eur. J. Appl. Physiol.*, 79 (6), 491-494.
- Riechman, S. E., Zoeller, R. F., Balasekaran, G. et al. (2002). Prediction of 2000 m indoor rowing performance using a 30 s sprint and maximal oxygen uptake. *J. Sports Sci.*, 20 (9), 681-687.
- Roth, W., Hasart, E., Wolf, W. & Pansold, P. (1983). Untersuchung zur Dynamik der Energiebereitstellung während maximaler Mittelzeitausdauerbelastung. *Med. Sport*, 23 (4), 107-114.
- Scott, B. R., Loenneke, J. P., Slattery, K. M. & Dascombe, B. J. (2016). Blood flow restricted exercise for athletes: A review of available evidence. *J. Sci. Med. Sport*, 19 (5), 360-367.
- Scott, N. W., McPherson, G. C., Ramsay, C. R. & Campbell, M. K. (2002). The method of minimization for allocation to clinical trials. a review. *Control Clin. Trials*, 23 (6), 662-674.
- Seiler, S. (2010). What is best practice for training intensity and duration distribution in endurance athletes? *Int. J. Sports Physiol. Perform.*, 5 (3), 276-291.
- Slysz, J., Stultz, J. & Burr, J. F. (2016). The efficacy of blood flow restricted exercise: A systematic review and meta-analysis. *J. Sci. Med. Sport*, 19 (8), 669-675.
- Steinacker, J. M., Lormes, W., Lehmann, M. & Altenburg, D. (1998). Training of rowers before world championships. *Med. Sci. Sports Exerc.*, 30 (7), 1158-1163.
- Suga, T., Okita, K., Morita, N. et al. (2009). Intramuscular metabolism during low-intensity resistance exercise with blood flow restriction. *J. Appl. Physiol.*, 106 (4), 1119-1124.
- Treff, G., Winkert, K., Machus, K. & Steinacker, J. M. (2018). Computer-aided stroke-by-stroke visualization of actual and target power allows for continuously increasing ramp tests on wind-braked rowing ergometers. *Int. J. Sports Physiol. Perform.*, 13 (6), 729-734.
- Treff, G., Winkert, K., Sareban, M. et al. (2017). Eleven-week preparation involving polarized intensity distribution is not superior to pyramidal distribution in national elite rowers. *Front. Physiol.*, 8: 515 (doi:10.3389/fphys.2017.00515).
- Vickers, A. J. & Altman, D. G. (2001). Statistics notes: Analysing controlled trials with baseline and follow up measurements. *BMJ*, 323 (7321), 1123 f.
- Wilson, J. M., Lowery, R. P., Joy, J. M. et al. (2013). Practical blood flow restriction training increases acute determinants of hypertrophy without increasing indices of muscle damage. *J. Strength Cond. Res.*, 27 (11), 3068-3075.

Korrespondenzadresse

Steffen Held, Abteilung für trainingswissenschaftliche Interventionsforschung, Deutsche Sporthochschule Köln, Am Sportpark Müngersdorf 6, 50933 Köln
E-Mail: s.held@dshs-koeln.de