

Billy Sperlich

INTENSITÄTSVERTEILUNG IM AUSDAUERTRAINING

Aktuelle Datenlage und Perspektiven

Literatur

Beaver, W. L., Wasserman K. & Whipp, B. J. (1986). A new method for detecting anaerobic threshold by gas exchange. *J. Appl. Physiol.*, 60 (6), 2020-2027.

Borg, G. (1970). Perceived exertion as an indicator of somatic stress. *Scand. J. Rehabil. Med.*, 2 (2), 92-98.

Bourgois, J., Steyaert, A. & Boone, J. (2013). Physiological and anthropometric progression in an international oarsman: a 15 year case study. *Int. J. Sports Physiol. Perform.*, 9 (4), 723-726.

Carnes, A. J. & Mahoney, S. E. (2018). Polarized vs. high intensity multimodal training in recreational runners. *Int. J. Sports Physiol. Perform.*: 1-28 (doi: 10.1123/ijsp.2018-0040).

Coffey, V. G. & Hawley, J. A. (2007). The molecular bases of training adaptation. *Sports Med.*, 37 (9), 737-763.

Conconi, F. et al. (1982). Determination of the anaerobic threshold by a noninvasive field test in runners. *J. Appl. Physiol.*, 52 (4), 869-873.

Coyle, E. F. et al. (1991). Physiological and biomechanical factors associated with elite endurance cycling performance. *Med. Sci. Sports Exerc.*, 23 (1), 93-107.

Davis, J. A. et al. (1976). Anaerobic threshold and maximal aerobic power for three modes of exercise. *J. Appl. Physiol.*, 41 (4), 544-550.

Dekerle, J., Vanhatalo, A. & Burnley, A. (2008). Determination of critical power from a single test. *Science & Sports*, 23 (5), 231-238.

Fiskerstrand, A. & Seiler, K. S. (2004). Training and performance characteristics among Norwegian international rowers 1970-2001. *Scand. J. Med. Sci. Sports*, 14 (5), 303-310.

Guellich, A., Seiler, S. & Emrich, E. (2009). Training methods and intensity distribution of young world-class rowers. *Int. J. Sports Physiol. Perform.*, 4 (4), 448-460.

Ingham, S. A. et al. (2008). Physiological and performance effects of low- versus mixed-intensity rowing training. *Med. Sci. Sports Exerc.*, 40 (3), 579-584.

Kindermann, W., Simon, G. & Keul, J. (1979). The significance of the aerobic-anaerobic transition for the determination of work load intensities during endurance training. *Eur. J. Appl. Physiol. Occup. Physiol.*, 42 (1), 25-34.

Mader, A. & Heck, H. (1986). A theory of the metabolic origin of "anaerobic threshold". *Int. J. Sports Med.*, 7 (Suppl. 1), 45-65.

Neal, C. M. et al. (2013). Six weeks of a polarized training-intensity distribution leads to greater physiological and performance adaptations than a threshold model in trained cyclists. *J. Appl. Physiol.*, 114 (4), 461-471.

Orie, J. et al. (2014). Thirty-eight years of training distribution in Olympic speed skaters. *Int. J. Sports Physiol. Perform.*, 9 (1), 93-99.

Plews, D. J. & Laursen, P. B. (2017). Training intensity distribution over a four-year cycle in Olympic Champion rowers: Different roads lead to Rio. *Int. J.*

Sports Physiol. Perform.: 1-24 (doi: 10.1123/ijsp.2017-0343).

Plews, D. J. et al. (2014). Heart-rate variability and training-intensity distribution in elite rowers. *Int. J. Sports Physiol. Perform.*, 9 (6), 1026-1032.

Riganas, C. S. et al. (2015). The rate of lactate removal after maximal exercise: the effect of intensity during active recovery. *J. Sports Med. Phys. Fitness*, 55 (10), 1058-1063.

Ronnby, S. et al. (2018). mHealth Self-Report Monitoring in competitive middle- and long-distance runners: Qualitative study of long-term use intentions using the technology acceptance model. *JMIR Mhealth Uhealth*, 6 (8): e10270 (doi: 10.2196/10270).

Scharhag-Rosenberger, F. et al. (2010). Exercise at given percentages of VO_2max : heterogeneous metabolic responses between individuals. *J. Sci. Med. Sport*, 13 (1), 74-79.

Seiler, S. (2010). What is best practice for training intensity and duration distribution in endurance athletes? *Int. J. Sports Physiol. Perform.*, 5 (3), 276-291.

Seiler, K. S. & Kjerland, G. O. (2006). Quantifying training intensity distribution in elite endurance athletes: is there evidence for an "optimal" distribution? *Scand. J. Med. Sci. Sports*, 16 (1), 49-56.

Sjodin, B. & Jacobs, I. (1981). Onset of blood lactate accumulation and marathon running performance. *Int. J. Sports Med.*, 2 (1), 23-26.

Sperlich, B., Becker, M., Hotho, A. et al. (2017). Sedentary behavior among national elite rowers during off-training - A pilot study. *Front. Physiol.*, 8: 655 (doi: 10.3389/fphys.2017.00655).

Sperlich, B. & Holmberg, H. C. (2017). The responses of elite athletes to exercise: An all-day, 24-h integrative view is required! *Front. Physiol.*, 8: 564 (doi: 10.3389/fphys.2017.00564).

Stegmann, H., Kindermann, W. & Schnabel, A. (1981). Lactate kinetics and individual anaerobic threshold. *Int. J. Sports Med.*, 2 (3), 160-165.

Stöggl, T. L. & Sperlich, B. (2015). The training intensity distribution among well-trained and elite endurance athletes. *Front. Physiol.*, 6: 295 (doi: 10.3389/fphys.2015.00295).

Stöggl, T. & Sperlich, B. (2014). Polarized training has greater impact on key endurance variables than threshold, high intensity, or high volume training. *Front. Physiol.*, 5: 33 (doi: 10.3389/fphys.2014.00033).

Treff, G. et al. (2017). Eleven-week preparation involving polarized intensity distribution is not superior to pyramidal distribution in national elite rowers. *Front. Physiol.*, 8: 515 (doi: 10.3389/fphys.2017.00515).

Treff, G. et al. (2019). The polarization-index: A simple calculation to distinguish polarized from non-polarized training intensity distributions. *Front. Physiol.*, 10: 707 (doi: 10.3389/fphys.2019.00707).

Urhausen, A. et al. (1993). Individual anaerobic threshold and maximum lactate steady state. *Int. J. Sports Med.*, 14 (3), 134-139.

Wasserman, K. & Mallroy, M. B. (1964). Detecting the threshold of anaerobic metabolism in cardiac patients during exercise. *Am. J. Cardiol.*, 14, 844-852.

Korrespondenzadresse

Univ.-Prof. Dr. Billy Sperlich, Integrative & Experimental Exercise Science & Training, Julius-Maximilians-Universität Würzburg, Judenbühlweg 11, 97082 Würzburg
E-Mail: billy.sperlich@uni-wuerzburg.de