

Kuno Hottenrott/Martin Möhle/Laura Hottenrott

KONTINUIERLICHE GLUKOSE-MESSUNG IM SPORT

Literatur

Boyne, M. S., Silver, D. M., Kaplan, J. & Saudek, C. D. (2003). Timing of changes in interstitial and venous blood glucose measured with a continuous subcutaneous glucose sensor. *Diabetes*, 52 (11), 2790-2794.

Davey, R. J., Low, C., Jones, T. W. & Fournier, P. A. (2010). Contribution of an intrinsic lag of continuous glucose monitoring systems to differences in measured and actual glucose concentrations changing at variable rates in vitro. *Journal of Diabetes Science and Technology*, 4 (6), 1393-1399.

Dickhuth, H.-H., Yin, L., Niess, A., Rucker, K., Mayer, F., Heitkamp, H.-C., Horstmann, T. (1990). Ventilatory, lactate-derived and catecholamine thresholds during incremental treadmill running: relationship and reproducibility. *International Journal of Sports Medicine*, 20 (2), 122-127.

Fayolle, C., Brun, J. F., Bringer, J., Mercier, J. & Renard, E. (2006). Accuracy of continuous subcutaneous glucose monitoring with the GlucoDay® in type 1 diabetic patients treated by subcutaneous insulin infusion during exercise of low versus high intensity. *Diabetes & Metabolism*, 32 (4), 313-320.

Iscoe, K. E., Campbell, J. E., Jamnik, V., Perkins, B. A. & Riddell, M. C. (2006). Efficacy of continuous real-time blood glucose monitoring during and after prolonged high-intensity cycling exercise: spinning with a continuous glucose monitoring system. *Diabetes Technology & Therapeutics*, 8 (6), 627-635.

Jansson, P. A., Fowelin, J., Smith, U. & Lönnroth, P. (1988). Characterization by microdialysis of intracellular glucose level in subcutaneous tissue in humans. *American Journal of Physiology*, 255 (2 Pt 1), E218-220.

Kropff, J., Bruttomesso, D., Doll, W., Farret, A., Galasso, S., Luijck, Y. M. et al. (2015). Accuracy of two continuous glucose monitoring systems: a head to head comparison under clinical research centre and daily life conditions. *Diabetes, Obesity and Metabolism*, 17 (4), 343-349.

Marunaka, Y. (2021). Roles of interstitial fluid pH and weak organic acids in development and amelioration of insulin resistance. *Biochemical Society Transactions*, 49 (2), 715-726.

Reiterer, F., Polterauer, P., Schoemaker, M., Schmelzeisen-Redecker, G., Freckmann, G., Heinemann, L. & Del Re, L. (2017). Significance and reliability of MARD for the accuracy of CGM systems. *Journal of Diabetes Science and Technology*, 11 (1), 59-67.

Sjöstrand, M., Gudbjörnsdóttir, S., Holmäng, A., Lönn, L., Strindberg, L. & Lönnroth, P. (2002). Delayed transcapillary transport of insulin to muscle interstitial fluid in obese subjects. *Diabetes*, 51 (9), 2742-2748.

Steil, G. M., Rebrin, K., Hariri, F., Jinagonda, S., Tardos, S., Darwin, C. & Saad, M. F. (2005). Interstitial fluid glucose dynamics during insulin-induced hypoglycaemia. *Diabetologia*, 48 (9), 1833-1840.

Weitgasser, R., Wielinger, H. & Pfohl, M. (2006). Stoffwechsellkontrolle: Glukosemessung, Bestimmung von Ketonkörpern, HbA1c und Fruktosamin. In: H. Schatz (Hrsg.), *Diabetologie kompakt – Grundlagen und Praxis* (S. 26-31). Stuttgart: Georg Thieme-Verlag.

Wilson, D. M., Beck, R. W., Tamborlane, W. V., Donchev, M. J., Kollman, C., Chase, P. et al. (2007). The accuracy of the FreeStyle Navigator continuous glucose monitoring system in children with type 1 diabetes. *Diabetes Care*, 30 (1), 59-64.

Yardley, J. E., Sigal, R. J., Kenny, G. P., Riddell, M. C., Lovblom, L. E. & Perkins, B. A. (2013). Point accuracy of interstitial continuous glucose monitoring during exercise in type 1 diabetes. *Diabetes Technology & Therapeutics*, 15 (1), 46-49.

Korrespondenzadresse

Univ.-Prof. Dr. Kuno Hottenrott, Arbeitsbereich Trainingswissenschaft und Sportmedizin, Institut für Sportwissenschaft, Martin-Luther-Universität Halle-Wittenberg, Von-Seckendorff-Platz 2, 06120 Halle (Saale)
E-Mail: kuno.hottenrott@sport.uni-halle.de